

CLAIMS

1. A crystallization method of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine methyl ester crystals exhibiting the specific peaks of diffracted X-rays at angles of diffraction (2θ , CuK α rays) of at least 6.0°, 24.8°, 8.2° and 16.5°, which comprises using water alone or a mixture of water and a lower alcohol as the crystallization solvent and maintaining the nucleation temperature at 30°C or greater.

2. The crystallization method of Claim 1, wherein said lower alcohol is methanol, and the methanol content in the whole solution to be subjected to crystallization is 15 wt.% or less.

3. A crystallization method of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine methyl ester crystals, which comprises using water alone or a mixture of water and a lower alcohol as the crystallization solvent and using, as the seed crystals, N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine methyl ester crystals exhibiting the specific peaks of diffracted X-rays at angles of diffraction (2θ , CuK α rays) of at least 6.0°, 24.8°, 8.2° and 16.5°, whereby the same type of crystals as the seed crystals are allowed to be preferentially precipitated.

4. The crystallization method of Claim 3, wherein said lower alcohol is methanol.

add
A2